

Bloch's conjecture revisited

by L.BARBIERI-VIALE and V.SRINIVAS

March '95

Abstract

Let X be a non-singular projective complex surface. We can show that Bloch's conjecture (*i.e.*, that if $p_g = 0$ then the Albanese kernel vanishes) is equivalent to the following statement:

If $p_g(X) = 0$ then for any given Zariski open $U \subset X$ and $\omega \in H^2(U, \mathbf{C})$ there is a smaller Zariski open $V \subset U$ such that

$$\omega|_V = \omega' + \omega_{\mathbf{Z}}$$

where $\omega' \in F^2 H^2(V, \mathbf{C})$ and $\omega_{\mathbf{Z}}$ is integral.

Let X denote a complex algebraic surface which is smooth and complete. Let $A_0(X)$ denote the subgroup of the Chow group of zero cycles given by cycles of degree zero, and let $J^2(X)$ be the Albanese variety of X , which we regard as the intermediate Jacobian canonically associated with the Hodge structure on $H^3(X, \mathbf{Z}(2))$. Let $\phi : A_0(X) \rightarrow J^2(X)$ be induced by the canonical (surjective) mapping to the Albanese variety. The kernel of ϕ is usually called the "Albanese kernel".

Bloch's conjecture *If $p_g(X) = 0$ then the "Albanese kernel" vanishes.*

The goal of this note is to show the following:

Theorem 1 *Let X be as above, and assume that $p_g(X) = 0$. Then*

$$\text{"Albanese kernel"} \cong \lim_{U \subset X} \frac{H^2(U, \mathbf{C})}{F^2 H^2(U, \mathbf{C}) + H^2(U, \mathbf{Z}(2))}$$

where the limit is taken over all non-empty Zariski open subsets of X .

We then clearly have the following:

Corollary 1 *Let X be as above, and assume that $p_g(X) = 0$. The "Albanese kernel" vanishes if and only if for any given Zariski open $U \subset X$ and $\omega \in H^2(U, \mathbf{C})$, there is a smaller Zariski open $V \subset U$ such that*

$$\omega|_V = \omega' + \omega_{\mathbf{Z}},$$

*where $\omega' \in F^2 H^2(V, \mathbf{C})$ and $\omega_{\mathbf{Z}}$ is integral (*i.e.*, $\omega_{\mathbf{Z}} \in \text{im } H^2(V, \mathbf{Z}(2))$).*

We will make use of the following result only in the particular case when Y is a curve.

Lemma 1 *Let Y be any reduced variety over the complex numbers. Then*

$$\lim_{U \subsetneq Y} \frac{H^1(U, \mathbf{C})}{F^1 H^1(U, \mathbf{C}) + H^1(U, \mathbf{Z}(1))} = 0$$

where the limit is taken over all non-empty Zariski open subsets of Y .

Proof Since the statement concerns the generic points of a variety, we may assume that Y is irreducible, non-singular and projective. For any open $U \subset Y$, let $S = Y - U$, and let $\text{Div}_S^0(Y)$ be the group of divisors on Y which are supported on S , and are homologous to 0 on Y . There is an exact sequence

$$0 \rightarrow H^1(Y, \mathbf{Z}(1)) \rightarrow H^1(U, \mathbf{Z}(1)) \xrightarrow{\text{res}} \text{Div}_S^0(Y) \rightarrow 0$$

where res denotes the sum of the residue maps associated to components of S which are divisors on Y . This underlies an exact sequence of mixed Hodge structures, where $F^1(\text{Div}_S^0(Y) \otimes \mathbf{C}) = \text{Div}_S^0(Y) \otimes \mathbf{C}$ (indeed, for any such divisor D , we have that

$$F^1 H_D^2(Y, \mathbf{C}(1)) \cong F^0 H^0(\tilde{D}, \mathbf{C}(0))$$

for a desingularization $\tilde{D}$ of D). Now

$$\frac{H^1(Y, \mathbf{C})}{F^1 H^1(Y, \mathbf{C})} \cong H^1(Y, \mathcal{O}_Y),$$

and since F^1 yields an exact functor on the category of mixed Hodge structures (see [D]), we get that

$$\frac{H^1(Y, \mathbf{C})}{F^1 H^1(Y, \mathbf{C})} \cong \frac{H^1(U, \mathbf{C})}{F^1 H^1(U, \mathbf{C})}.$$

Hence there is a natural map

$$\text{Div}_S^0(Y) \rightarrow \frac{H^1(U, \mathbf{C})}{F^1 H^1(U, \mathbf{C}) + H^1(Y, \mathbf{Z}(1))} \cong \frac{H^1(Y, \mathbf{C})}{F^1 H^1(Y, \mathbf{C}) + H^1(Y, \mathbf{Z}(1))} \cong \text{Pic}^0(Y).$$

This map is known to be just the natural map $D \mapsto \mathcal{O}_Y(D)$. Hence

$$\frac{H^1(U, \mathbf{C})}{F^1 H^1(U, \mathbf{C}) + H^1(U, \mathbf{Z}(1))} \cong \frac{\text{Pic}^0(Y)}{\text{subgroup generated by } \text{Div}_S^0(Y)}.$$

Since $\varinjlim_S \text{Div}_S^0(Y)$ is the group of all divisors on Y which are homologous to 0, which maps surjectively to $\text{Pic}^0(Y)$, the Lemma follows immediately. •

Remark: Let Y be a non-singular projective complex variety. By the same argument in the proof of the Lemma, one can see that

$$\lim_{U \subsetneq Y} \frac{H^{2p-1}(U, \mathbf{C})}{F^p H^{2p-1}(U, \mathbf{C}) + H^{2p-1}(U, \mathbf{Z}(p))} = 0$$

(where now the limit is taken over Zariski open subsets U of Y such that the codimension of $Y - U$ is $\geq p$), if and only if $J^p(Y)$ is algebraic *i.e.*, the Abel-Jacobi map is surjective.

Proof of the Theorem Let $\mathcal{H}^2(\mathbf{Z}(2))$ and $\mathcal{H}^2/\mathcal{F}^2$ be the Zariski sheaves on X associated to $U \mapsto H^2(U, \mathbf{Z}(2))$ and $U \mapsto H^2(U, \mathbf{C})/F^2$ respectively. Let $\mathcal{H}^3(\mathbf{Z}(2)_{\mathcal{D}})$ denote the Deligne–Beilinson cohomology Zariski sheaf on X (*cf.* [G] and [E]). We then have the following diagram of sheaves

$$\begin{array}{ccccccc}
0 & \rightarrow & \mathcal{H}^2(\mathbf{Z}(2)) & \rightarrow & H^2(\mathbf{C}(X), \mathbf{Z}(2)) & \rightarrow & \coprod_{x \in X^1} i_x H^1(\mathbf{C}(x), \mathbf{Z}(1)) \rightarrow \dots \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & \mathcal{H}^2/\mathcal{F}^2 & \rightarrow & H^2(\mathbf{C}(X))/F^2 & \rightarrow & \coprod_{x \in X^1} i_x H^1(\mathbf{C}(x))/F^1 \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & \mathcal{H}^3(\mathbf{Z}(2)_{\mathcal{D}}) & \rightarrow & H^3(\mathbf{C}(X), \mathbf{Z}(2)_{\mathcal{D}}) & \rightarrow & \coprod_{x \in X^1} i_x H^2(\mathbf{C}(x), \mathbf{Z}(1)_{\mathcal{D}}) \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

obtained by comparing the arithmetic resolutions of the left hand column (see [BV] for that of $\mathcal{H}^2/\mathcal{F}^2$) via the standard long exact sequences in Deligne–Beilinson cohomology (and taking into account that singular cohomology of an affine vanishes in degrees greater than its algebraic dimension). Now, because of the Lemma, we have that

$$H^2(\mathbf{C}(x), \mathbf{Z}(1)_{\mathcal{D}}) \stackrel{\text{def}}{=} \varinjlim_{U \subset \overline{\{x\}}} H^2(U, \mathbf{Z}(1)_{\mathcal{D}}) \cong \varinjlim_{U \subset \overline{\{x\}}} \frac{H^1(U, \mathbf{C})}{F^1 H^1(U, \mathbf{C}) + H^1(U, \mathbf{Z}(1))} = 0.$$

Thus the right-bottom corner in the diagram above vanishes and the sheaf $\mathcal{H}^3(\mathbf{Z}(2)_{\mathcal{D}})$ is then identified by the constant sheaf associated with the group

$$H^3(\mathbf{C}(X), \mathbf{Z}(2)_{\mathcal{D}}) \stackrel{\text{def}}{=} \varinjlim_{U \subset X} H^3(U, \mathbf{Z}(2)_{\mathcal{D}}) \cong \varinjlim_{U \subset X} \frac{H^2(U, \mathbf{C})}{F^2 H^2(U, \mathbf{C}) + H^2(U, \mathbf{Z}(2))}$$

where the limit is taken over all non-empty Zariski open subsets of X . Finally, if $p_g(X) = 0$, it is known (see [BV] and [E]) that

$$H^0(X, \mathcal{H}^3(\mathbf{Z}(2)_{\mathcal{D}})) \cong \ker(A_0(X) \xrightarrow{\phi} J^2(X)),$$

yielding the claimed identification. •

Remark: Note that for a surface X with $p_g \neq 0$ we have that

$$\varinjlim_{U \subset X} \frac{H^2(U, \mathbf{C})}{F^2 H^2(U, \mathbf{C}) + H^2(U, \mathbf{Z}(2))} \neq 0,$$

since this group maps onto the “Albanese kernel”, which is non-zero by [M].

For a surface X with $p_g = 0$, the canonical “residue map”

$$H^2(\mathbf{C}(X))/F^2 \rightarrow \coprod_{x \in X^1} H^1(\mathbf{C}(x))/F^1$$

is injective, with quotient group isomorphic to $H^3(X, \mathbf{C})/F^2$ (see [BV]); thus the integral lifting of the “residue” of a given $\omega \in H^2(\mathbf{C}(X))/F^2$ always exists, but, in order to lift ω itself to a class in $H^2(\mathbf{C}(X), \mathbf{Z}(2))$, one should know that the lifting of the “residue” of ω yields zero in the group of zero-cycles as well as in $H^3(X, \mathbf{Z}(2))$.

References

- [BV] L.BARBIERI-VIALE: On the Deligne–Beilinson cohomology sheaves, preprint, 1994.
- [D] P.DELIGNE: Théorie de Hodge II,III , *Publ. Math. IHES* **40** (1972) 5–57 and **44** (1974) 5–78.
- [E] H.ESNAULT: A note on the cycle map, *J.reine angew.Math.* **411** (1990) 51–65.
- [G] H.GILLET: Deligne homology and the Abel–Jacobi maps, *Bull.AMS* **10** (1984) 285–288.
- [M] D.MUMFORD: Rational equivalence of 0-cycles on surfaces, *J.Math.Kyoto Univ.* **9** (1968) 195–204.

Dipartimento di Matematica
Università di Genova
Via L.B.Alberti, 4
16132 – GENOVA
Italia

Tata Institute of Fundamental Research
School of Mathematics
Homi Bhabha Road
400 005 – BOMBAY
India